RESEARCH PROTOCOL

Harnessing RNA Interference to Constrain Ustilago hordei Growth and Barley Infection

Emily Jiang, BSc Candidate [1]*, Olivia Xu, BMSc Candidate [2]

 Department of Neuroscience, The University of California, Los Angeles, Los Angeles, California, United States, 90025
Department of Biochemistry, Schulich School of Medicine & Dentistry, Western University, London, Ontario, Canada, N6A 5C1

*Corresponding Author: emilyjiang003@ucla.edu

Abstract

Introduction: Ustilago hordei (U. hordei), a facultative biotrophic fungal pathogen, poses a significant threat to barley (Hordeum vulgare) crops, causing barley smut and substantial economic losses. Fungal colonies have a high risk of developing resistance to traditional fungicides upon usage and can pose potential risks to the environment. The survival and infection cycle of U. hordei is enabled through a number of genes suppressing the host immune system, including UhAvr1, Fly1, and the beta-glucuronidase gene (GUS). This research project aims to explore the application of RNA interference (RNAi) to silence these genes implicated in barley colonization, aiming to understand the effect of Spray-Induced Gene Silencing (SIGS) using double-stranded RNA (dsRNA) on U. hordei growth.

Methods: Three pivotal genes (*UhAvr1, Fly1, GUS*) important to *U. hordei* infectivity were chosen to be amplified and transformed into dsRNA that contains small interfering RNA (siRNA) regions through in vitro expression in *Escherichia coli*. These genes were selected due to the roles they play in promoting *U. hordei* virulence and enhancing *U. hordei's* ability to metabolize complex carbohydrates. The impact of dsRNA on *U. hordei* growth will be assessed at concentrations of 50 ng/µL, 100 ng/µL, 150 ng/µL, 200 ng/µL. After growth is assessed, infectivity will be measured by comparing the number of barley plants infected versus the number of plants inoculated 6 days after inoculation.

Results: The application of target gene dsRNA will limit fungal growth and barley colonization. A positive relationship between translational inhibition and dsRNA concentration was observed in *U. hordei* populations. Furthermore, there was a negative relationship between the concentration of dsRNA applied to *U. hordei* and the fraction of plants infected versus inoculated.

Conclusion: Ultimately, this study provides insights into the application of the three potential dsRNAs in SIGS as a replacement for chemical fungicides with wide agricultural applications.

Keywords: ustilago hordei; barley smut; RNA interference; fungicide

Introduction

Ustilago hordei (U. hordei) is a facultative biotrophic fungal pathogen that primarily affects Hordeum vulgare, a worldwide economically and nutritionally important crop colloquially known as barley, causing a disease known as barley smut [1]. Barley smut is a common fungal disease that can lead to reduced yield and quality of barley crops and significant economic losses. U. hordei has a complex life cycle that involves both sexual and asexual stages that are coupled with its pathology [2]. These resting spores of U. hordei, named teliospores, act as the main markers of massive proliferation and are usually dispersed onto healthy barley, beginning the infection cycle [3, 4]. Currently, methods of disease management include crop rotation, using disease-resistant barley varieties, and fungicides [5]. However, the pollution resulting from fungicide usage and the preferential selection for fungicide-resistant U. hordei

Jiang et al | URNCST Journal (2024): Volume 8, Issue 10 DOI Link: <u>https://doi.org/10.26685/urncst.651</u> can pose serious long-lasting problems to the environment [5]. Recently, an RNA silencing approach known as Spray-Induced Gene Silencing (SIGS), which utilizes the nature of fungi to intake dsRNA upon exposure, provides a novel approach to combat fungal pathogens [6].

RNA interference (RNAi) is a natural sequence-specific gene regulation process using small RNA molecules and has become increasingly important as a tool for research in plants. The initiation of RNAi involves the cleavage of trigger RNA (dsRNA) into short interfering RNAs (siRNA) by the enzyme, Dicer. The siRNAs interact with the enzyme Argonaute (AGO) and are unwound into 2 single-stranded RNAs called sense and antisense RNA. The sense RNA is degraded by AGO, while the antisense RNA is incorporated into the RNA-inducing silencing complex (RISC), which allows the antisense RNA to hybridize with its complementary target mRNA [7]. This will lead to the



OPEN ACCESS

cleavage of target mRNA, resulting in target gene silencing. This research project aims to use RNAi to silence genes that are involved during colonization of barley and evaluate the effect of this treatment on *U. hordei* growth [8].

Several genes crucial to U. hordei pathogenicity were selected in this research project. Three genes, UhAvr1, Fly1, and the beta-glucuronidase gene (GUS) were chosen based on their functions and expression during barley infection. UhAvr1 was selected as it actively expresses itself in the early stages of infection, leading to a general suppression in system immune the host [9]. The Flv1 gene is present in both U. hordei and Ustilago maydis, the perpetrator of corn smut. It codes for fungalysin metalloprotease, which reduces plant chitinase activity [10]. Acting as the first line of defense, plant chitinase restricts fungal growth through fungal cell wall remodeling [11]. The expression of fungalysin metalloprotease, as a result, tears a hole in the airtight host defense mechanism, creating room for U. hordei survival. The third gene on the list, GUS, codes for the production of β -glucuronidase, a lysosomal enzyme responsible for breaking down complex carbohydrates in the host, influencing the structure of the barley cell wall [12]. Therefore, the GUS gene is also known to play an important role in barley colonization, and its product, β -glucuronidase, is produced in the tips of the hyphae of Ustilago hordei in infected barley tissue [13].

We hypothesize that the utilization of RNA interference with genes *UhAvr1*, *Fly1*, *GUS* is an effective mechanism in restricting the growth of *Ustilago hordei* and could potentially limit *U. hordei's* growth and ability to infect barley plants.

Methods

Culture Maintenance and Creating cDNA Library

The cultivation of U. hordei will be conducted in YEPS_{light} (0.4% yeast extract, 0.4% peptone, and 2% saccharose). The culture will be maintained in liquid medium at 22 °C with continuous shaking at 200 rpm until the optimal density 600nm (OD₆₀₀) reaches 1.0 [1]. To produce a complementary DNA (cDNA) library for U. hordei, RNA will be extracted using a MasterpureTM Complete DNA and RNA Purification Kit (Epicentre®, Illumina®, MC89010). Genomic DNA is removed by the addition of ezDNase enzyme for 2 min at 37 °C. ezDNase is inactivated under 55 °C. To create cDNA strands using the RNA template, first mix 0.01 pg-0.5 µg template RNA with 0.5-1.0 µM of gene-specific primers and 0.5-1.0 mM of dNTPs. Add buffer (DTT+RNase inhibitor) and nucleasefree water to the reaction as well. When the mix is ready, add 1 µL of avian myeloblastosis virus (AMV) reverse transcriptase with 5 x AMV RT buffer to the solution; put the reaction mix under 42 °C for 60 minutes. To inactivate the enzyme, raise the temperature of the reaction mix to 70-85 °C for 5 to 15 min [14]. The cDNA products contain the U. hordei target genes that are essential for the colonization of barley that will be amplified using polymerase chain

Jiang et al | URNCST Journal (2024): Volume 8, Issue 10 DOI Link: <u>https://doi.org/10.26685/urncst.651</u> reaction (PCR) and used as the template for the synthesis of dsRNA in later steps [15].

Choosing the siRNA for the Genes of Interest

The sidirect database will be used in the selection of suitable siRNA for the target genes *UhAvr1*, *Fly1*, *GUS*. siRNAs are often 20-30 nucleotides in length [16]. The database can identify the effective regions for siRNA of target genes. For our project, we will identify three effective areas for each gene. These regions help build effective siRNA and prevent excessive off-target trial-and-error [17].

dsRNA Synthesis

After the creation of the cDNA library for Ustilago hordei and the selection of target genes, PCR and primers specific to the siRNA of UhAvr1, Flv1, and GUS are used to amplify the genes of interest. The target gene amplicons are purified and collected through gel elution using QIAquick Gel Extraction Kit (QIAGEN, 28704). This project will use engineered bacterial expression to produce dsRNA. The bacterial strain selected is Escherichia. coli (E. coli) HT115 due to its familiar genetic background, ease of genetic manipulation, and lack of RNase III. A lack of RNase III is important to prevent the degradation of dsRNA, our final product. The vector we chose is L4440 with oppositely oriented T7 promoters, a DNA sequence that allows T7 RNA polymerase (T7 RNAP) to recognize and initiate transcription, due to its reliable presence in literature and widespread use in other studies. T7 RNAP will be synthesized on induction by isopropyl β -D-1thiogalactopyranoside (IPTG), which will synthesize complementary strands of RNA, forming dsRNA [18]. Gel electrophoresis will be used to confirm the presence of dsRNA. A band of the corresponding size will be observed if our target dsRNA is on the gel. The product is purified using ion pair reverse-phase high-performance liquid chromatography [19].

Testing the Growth Rate and Infectivity After dsRNA Intake

Following the extraction of the dsRNA from E. coli, the liquid culture medium of U. hordei will be treated at 50 ng/µL, 100 ng/µL, 150 ng/µL, 200 ng/µL of dsRNA [16]. Corresponding control groups (SS20 Buffer of the same volume only) for each treatment group will also be included in the experimental setup. This project will incubate the culture for 12 hrs and 24 hrs, then measure the OD_{600} of U. hordei in distilled water to quantify U. hordei growth [1]. We will use 1:1 solution of distilled water and YEPS_{light} as a blank to which other values will be compared relatively. If there is the presence of viable U. hordei after dsRNA treatment, these colonies will be exposed to barley to test for infectivity. Pathogenicity was measured by comparing the number of barley plants infected versus the number of plants inoculated 6 days after inoculation. Infected plants will show visible black spots on the barley seeds [20].

Results

This project expects that cell culture will produce viable U. hordei fungal colonies that express the target gene proteins (UhAcr1, Fly1, GUS). A complete cDNA library is expected to be created based on the transcriptome of U. hordei. The PCR on cDNA for the three target genes will amplify the target gene sequence substantially. These cDNA fragments act as the precursor for the production of dsRNA, which contains the potential siRNA effector sequences obtained from the sidirect database search. This project expects the target gene dsRNA product, when applied to the U. hordei culture, to limit fungal growth and colonization of barley plants. More specifically, it is expected that there will be a positive relationship between the effect of translational inhibition on U. hordei exhibited by dsRNA treatment and the concentration of dsRNA. The U. hordei, if viable colonies exist after dsRNA treatment, is anticipated to decrease its infectivity. In other words, the higher the concentration of dsRNA, the lower the growth of U. hordei in liquid medium, and fewer infected barley plants will be observed.

Conclusions

This research project provides new insights into RNAi as an alternative to chemical fungicides and is a potential antifungal method in the face of rising antifungal resistance. Using SIGS, dsRNA of the three genes that we identified (*UhAcr1, Fly1, GUS*), can be applied in a convenient manner. Therefore, considering these expected results, our research can contribute to the field of agriculture by identifying alternative or supporting solutions to chemical fungicides.

List of Abbreviations Used

GUS: *Beta-glucuronidase gene* RNAi: RNA interference siRNA: Small interfering RNA dsRNA: double stranded RNA SIGS: spray-induced gene silencing AGO: argonaute RISC: RNA-induced silencing complex cDNA: complementary DNA OD: optimal density YEPS_{light}: 0.4% yeast extract, 0.4% peptone, and 2% saccharose solution PCR: polymerase chain reaction IPTG: isopropyl β-D-1-thiogalactopyranoside

Conflicts of Interest

The author(s) declare that they have no conflict of interests.

Ethics Approval and/or Participant Consent

No animals or human subjects were used or studied in the making of this research proposal project. Therefore, no ethics approval nor participant consent is necessary.

Jiang et al | URNCST Journal (2024): Volume 8, Issue 10 DOI Link: <u>https://doi.org/10.26685/urncst.651</u>

Authors' Contributions

EJ: made substantial contributions to the design of the study, reviewed literature, drafted the manuscript, and gave final approval of the version to be published. Is in agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. OX: made substantial contributions to the design of the study, reviewed literature, drafted the manuscript, and gave final approval of the version to be published. Is in agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding

This study was not funded.

References

- Ökmen B, Schwammbach D, Bakkeren G, Neumann U, Doehlemann G. The Ustilago hordei–Barley Interaction is a Versatile System for Characterization of Fungal Effectors. JoF. 2021 Jan 27;7(2):86. <u>https://doi.org/10.3390/jof7020086</u>
- [2] Zuo W, Ökmen B, Depotter JRL, Ebert MK, Redkar A, Misas Villamil J, et al. Molecular Interactions Between Smut Fungi and Their Host Plants. Annu Rev Phytopathol. 2019 Aug 25;57(1):411–30. <u>https://doi.org/10.1146/annurev-phyto-082718-100139</u>
- [3] Hu GG, Linning R, Bakkeren G. Sporidial mating and infection process of the smut fungus, Ustilago hordei, in susceptible barley. Can J Bot. 2002 Oct 1;80(10):11 03–14. <u>https://doi.org/10.1139/b02-098</u>
- [4] Gaudet DA, Wang Y, Penniket C, Lu ZX, Bakkeren G, Laroche A. Morphological and Molecular Analyses of Host and Nonhost Interactions Involving Barley and Wheat and the Covered Smut Pathogen Ustilago hordei. MPMI. 2010 Dec;23(12):1619–34. <u>https://doi.org/10.1094/MPMI-11-09-0271</u>
- [5] Woldemichael MD. Importance, Biology, Epidemiology, and Management of Loose Smut (Ustilago nuda) of Barley (Hordeum vulgare): A Review. 13.
- [6] Székács A, Ammour AS, Mendelsohn ML. Editorial: RNAi Based Pesticides. Front Plant Sci. 2021 Jul 29; 12:714116. <u>https://doi.org/10.3389/fpls.2021.714116</u>
- [7] Koeppe S, Kawchuk L, Kalischuk M. RNA Interference Past and Future Applications in Plants. IJMS. 2023 Jun 5;24(11):9755. <u>https://doi.org/10.33</u> <u>90/ijms24119755</u>
- [8] Xu W, Jiang X, Huang L. RNA Interference Technology. In: Comprehensive Biotechnology [Internet]. Elsevier; 2019 [cited 2023 Nov 22]. p. 560– 75. Available from: <u>https://linkinghub.elsevier.com/</u> retrieve/pii/B9780444640468002822

- [9] Montenegro Alonso AP, Ali S, Song X, Linning R, Bakkeren G. UhAVR1, an HR-Triggering Avirulence Effector of Ustilago hordei, Is Secreted via the ER– Golgi Pathway, Localizes to the Cytosol of Barley Cells during in Planta-Expression, and Contributes to Virulence Early in Infection. JoF. 2020 Sep 18;6(3): 178. https://doi.org/10.3390/jof6030178
- [10] Ökmen B, Kemmerich B, Hilbig D, Wemhöner R, Aschenbroich J, Perrar A, et al. Dual function of a secreted fungalysin metalloprotease in Ustilago maydis. New Phytologist. 2018 Oct;220(1):249–61. https://doi.org/10.1111/nph.15265
- [11] Silipo A, Erbs G, Shinya T, Dow JM, Parrilli M, Lanzetta R, et al. Glyco-conjugates as elicitors or suppressors of plant innate immunity. Glycobiology. 2010 Apr 1;20(4):406–19. <u>https://doi.org/10.1093/glycob/cwp201</u>
- [12] Fior S, Gerola PD. Impact of ubiquitous inhibitors on the GUS gene reporter system: evidence from the model plants Arabidopsis, tobacco and rice and correction methods for quantitative assays of transgenic and endogenous GUS. Plant Methods. 2009;5(1):19. <u>https://doi.org/10.1186/1746-4811-2-19</u>
- [13] Jefferson RA, Kavanagh TA, Bevan MW. GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. The EMBO Journal. 1987 Dec;6(13):3901–7. <u>https://doi.org/10. 1002/j.1460-2075.1987.tb02730.x</u>
- [14] Reverse Transcription Setup | Thermo Fisher Scientific - UK [Internet]. Thermofisher.com. 2016 [cited 2024 Sept 07]. Available from: <u>https://www.thermofisher.com/us/en/home/life-science/cloning/cloning-learning-center/invitrogen-school-of-molecular-biology/rt-education/reverse-transcription-setup.html</u>

- [15] Parkinson J, editor. Expressed Sequence Tags (ESTs): Generation and Analysis [Internet]. Totowa, NJ: Humana Press; 2009 [cited 2023 Nov 22]. (Methods in Molecular Biology; vol. 533). Available from: <u>https://</u> link.springer.com/10.1007/978-1-60327-136-3
- [16] Dang Y, Yang Q, Xue Z, Liu Y. RNA Interference in Fungi: Pathways, Functions, and Applications. Eukaryot Cell. 2011 Sep;10(9):1148–55. <u>https://doi.org/10.1128/ec.05109-11</u>
- [17] Bhagta S, Bhardwaj V, Kant A. Exogenous dsRNA trigger RNAi in Venturia inaequalis resulting in down regulation of target genes and growth reduction. Mol Biol Rep. 2023 Oct;50(10):8421–9. <u>https://doi.org/10. 1007/s11033-023-08736-3</u>
- [18] Nwokeoji AO, Kung AW, Kilby PM, Portwood DE, Dickman MJ. Purification and characterisation of dsRNA using ion pair reverse phase chromatography and mass spectrometry. Journal of Chromatography A. 2017 Feb;1484:14–25. <u>https://doi.org/10.1016/j.chr</u> <u>oma.2016.12.062</u>
- [19] Guan R, Chu D, Han X, Miao X, Li H. Advances in the Development of Microbial Double-Stranded RNA Production Systems for Application of RNA Interference in Agricultural Pest Control. Front Bioeng Biotechnol. 2021 Sept 13;9:753790. <u>https://doi.org/ 10.3389/fbioe.2021.753790</u>
- [20] Ali S, Laurie JD, Linning R, Cervantes-Chávez JA, Gaudet D, Bakkeren G. An Immunity-Triggering Effector from the Barley Smut Fungus Ustilago hordei Resides in an Ustilaginaceae-Specific Cluster Bearing Signs of Transposable Element-Assisted Evolution. Xu JR, editor. PLoS Pathog. 2014 Jul 3;10(7):e1004223. <u>https://doi.org/10.1371/journal.ppat.1004223</u>

Article Information

Managing Editor: Jeremy Y. Ng Peer Reviewers: Alita Gideon, Mikayla Hall-Bruce Article Dates: Received Jun 11 24; Accepted Sep 07 24; Published Oct 02 24

Citation

Please cite this article as follows: Jiang E, Xu O. Harnessing RNA interference to constrain *ustilago hordei* growth and barley infection. URNCST Journal. 2024 Oct 02: 8(10). <u>https://urncst.com/index.php/urncst/article/view/651</u> DOI Link: <u>https://doi.org/10.26685/urncst.651</u>

Copyright

© Emily Jiang, Olivia Xu. (2024). Published first in the Undergraduate Research in Natural and Clinical Science and Technology (URNCST) Journal. This is an open access article distributed under the terms of the Creative Commons Attribution License (<u>https://creativecommons.org/licenses/by/4.0/</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Undergraduate Research in Natural and Clinical Science and Technology (URNCST) Journal, is properly cited. The complete bibliographic information, a link to the original publication on <u>http://www.urncst.com</u>, as well as this copyright and license information must be included.



Do you research in earnest? Submit your next undergraduate research article to the URNCST Journal! | Open Access | Peer-Reviewed | Rapid Turnaround Time | International | | Broad and Multidisciplinary | Indexed | Innovative | Social Media Promoted | Pre-submission inquiries? Send us an email at <u>info@urncst.com</u> | <u>Facebook</u>, <u>Twitter</u> and <u>LinkedIn</u>: @URNCST Submit YOUR manuscript today at <u>https://www.urncst.com</u>!